



RANGE EXTENSION OF TWO BRACHYURAN CRABS COLLECTED FROM THE AMAZON CONTINENTAL SHELF

Déborah Elena Galvão Martins^{1,2*} ; Flavio de Almeida Alves-Júnior^{1,2} ; Israel Hidenburgo Aniceto Cintra¹ ; Bianca Bentes²

Abstract

The Amazon Continental Shelf (ACS) is composed of numerous benthic habitats, which shelter a high diversity of invertebrates, especially crustaceans. However, due to the complexity of ecosystems, knowledge of crustacean fauna is still incomplete. We herein report the northernmost Brazilian record of two brachyuran crab species, *Teleophrys pococki* Rathbun, 1924 (Mithracidae) and *Euchirops americanus* A. Milne-Edwards, 1880 (Plagusidae), associated with the Great Amazon Reef System (GARS) (State of Amapá), additionally, providing a description of their first gonopods (G1). Both male specimens *E. americanus* and *T. pococki*, were accidentally collected during commercial fishing operation of the red snapper *Lutjanus purpureus* (Poey, 1866) in the ACS (northern Brazil), at depths up to 70 m, in August 2023 and September 2025. The presence of both species in the GARS increases crustacean biodiversity in this ecosystem and highlights the need for further studies in the region to assess its true diversity and to develop management plans focused on species conservation.

Keywords: Amazon Reefs. Bycatch. State of Amapá. Geographic distribution. New record.

¹Laboratório de Crustáceos (LabCrus), Socio-Environmental and Water Resources Institute (ISARH), Federal Rural University of Amazônia (UFRA), Avenida Presidente Tancredo Neves, nº 2501, Terra Firme, CEP: 66077-830, Belém, Pará, Brasil

²Núcleo de Ecologia Aquática e Pesca da Amazônia (NEAP), Post-Graduate Programme in Aquatic Ecology and Fisheries (PPGEAP), Federal University of Pará (UFPA), Avenida Perimetral, 2651, Terra Firme, 66077-530 Belém, Pará, Brasil

*Corresponding author: deborah.martins@ufra.edu.br

Submitted on: 17 Nov. 2025
Accepted on: 11 Dec. 2025
Published on: 20 Jan. 2026

1 Introduction

The Amazon continental shelf (ACS) is classified as a hotspot of diversity, due to the elevated complexity of habitats, the wide range of ecological interactions and adequate environmental conditions for the establishment of species (KLAUTAU et al., 2025); much of this biodiversity is due to the presence of the Amazon River plume, which transports organic matter, establishing a high energy flow in marine food chains (WEBER et al., 2017; DRAKE et al., 2021).

In this area, there is also an ecosystem composed of rhodoliths, and coral beds designated as the Great Amazon Reef System (GARS), which is distributed along mesophotic areas of ACS, ranging from 20 to ~100 m (VALE et al., 2022).

The GARS is the target of several studies, especially on geological mapping, ecological functionality and the diversity associated with this environment (MOURA et al., 2016; VALE et al., 2022).

However, this region has already been affected by human activities, such as fishing, marine litter and the effects of climate change (ALVES-JÚNIOR et al., 2024a), even before the knowledge about the species diversity (including crustaceans, which are usually recorded as part of fishing bycatch) was completely available (MARTINS; ALVES-JÚNIOR, CINTRA, 2025).

The family Mithracidae MacLeay, 1838 corresponds to 15 genera and 68 species, occurring from the coastal zones to the continental shelf break, associated with sand, gravel or coral bottoms (MELO, 1996; WINDSOR; FELDER, 2014; DECANET, 2025a).

Into this family, the genus *Teleophrys* Stimpson, 1860 comprises five recent species: *T. cristulipes* Stimpson, 1860, *T. ornatus* Rathbun, 1901, *T. pococki* Rathbun, 1924, *T. ruber* (Stimpson, 1871), *T. tumidus* (Cano, 1889), besides *T. acornis* Portell & Collins, 2004 as a fossil record (DECANET, 2025a).

From the Brazilian waters, only the species *T. ornatus* and *T. pococki* have been recorded in the northeastern and southeastern regions (MELO, 1996). Despite the occurrence of *T. pococki* in coastal zones of Brazil, this species was reported in a few localities, represented only by occasional records.

Additionally, crabs of the family Plagusiidae Dana, 1851 are found along the coastal zones and continental slope areas (including Oceanic Islands), associated with bottoms of sand, gravel and coral (MELO, 1996; 1998).

These organisms are widely reported along the Atlantic and Indo-Pacific Oceans, which are classified in 21 species distributed into seven genera: *Caligoplagusia* Fujita & Narusa, 2014, *Davusia* Guinot, 2007, *Euchirograpsus* H. Milne Edwards, 1853, *Guinusia* Schubart & Cuesta, 2010, *Miersiograpsus* Türkay, 1978 and *Plagusia* Latreille, 1804, as well as the extinct species *Petrusia* Beschin, Busulini, Tessier & Zorzin, 2016 (GUINOT, 2007; NG; GUINOT; DAVIE, 2008; DECANET, 2025b).

Currently, the genus *Euchirograpsus* is composed of nine valid species, occurring from 10 to 510 m of depth in the bottoms of sand, mud, rock and corals, associated with gorgonians and sponges (TÜRKAY, 1975; MANNING; HOLTHUIS, 1981, ALVES-JÚNIOR et al., 2016; WILSON; POHLE, 2016). In Brazilian waters, only two species have been recorded: *Euchirograpsus antillensis* Türkay, 1975 and *Euchirograpsus americanus* A. Milne-Edwards, 1880, both occurring along the continental shelf and shelf break of the northeastern and southeastern regions (MELO, 1996; 1998; ALVES-JÚNIOR et al., 2016). Thus, herein we report the Brazilian northernmost records of two brachyuran species: *Teleophrys pococki* and *Euchirograpsus americanus* from the state of Amapá, associated with the GARS.

2 Material and Methods

The crab species were obtained incidentally attached to an iron cage trap called "manzuá" (mesh size of 5 cm) during commercial fishing operations of the red snapper *Lutjanus purpureus* (Poey, 1866). These fishing activities were performed between the depths of 70 and 100 m, along the continental shelf of the Brazilian state of Amapá (Figure 1) from August 2023 to September 2025, covering areas above the Great Amazon Reef System (GARS).

In the field, the rhodoliths adhered to the external surface of the trap were manually removed, related invertebrate species were fixed in 70% ethanol, and then transported to the *Laboratório de Crustáceos* (LabCrus). All sampling procedures were supervised by the *Centro Nacional de Pesquisa e Conservação da Biodiversidade Marinha do Norte* (CEPNOR) under the SISBIO Number 44915-3.

In the laboratory, both species were identified following Türkay (1975) and Melo (1996), photographed, measured in carapace width (cw.), carapace length (cl.) and wet weight (ww.). After the procedures, the specimens were deposited under voucher number at the *Coleção Carcinológica do Laboratório de Crustáceos* (LabCrus) / *Universidade Federal Rural da Amazônia* (UFRA).




Figure 1. Map of the study area, covering the Amazon continental shelf (ACS), indicating the occurrence sites of *Teleophrys pococki* Rathbun, 1924 and *Euchirograpsus americanus* A. Milne-Edwards, 1880. Shapefile of the GARS from Moura et al. (2016).

3 Results and Discussion

One adult male of *T. pococki* (Figure 2 A-C) (cw. 11.6 mm, cl. 10.3 mm and ww. 0.57 g) (Voucher Number: 2.16.1A) was collected at a depth of 70 m in the GARS area (August 2023), associated with rhodoliths in the gravel bottom, state of Amapá (Brazil) (04°43'59" N; 050°39'18" W). According to Tavares and Mendonça Jr. (2022), the bathymetric distribution of *T. pococki* covers continental shelf areas, from shallow waters to 100 m; however, it is more commonly found up to 25 m of depth (MELO, 1996; CASTAÑO; CAMPOS, 2003; CARMONA-SUÁREZ; POUPIN, 2016).

Our specimen fits well with the original description provided by Rathbun (1924) and additional information provided by Rathbun (1925) and Melo (1996), which include the main characteristics: carapace larger than long, smooth, containing few granules or small tubercles which are more visible in the branchial and cardiac regions; rostrum bifid, also larger than long, with one antero-lateral teeth; carpus of chelipeds with two lobes only in the inner face, chelipeds smooth forming an open angle in the centre, with curved edges. Ambulatory legs with several tubercles forming crests on the upper surface (Figure 2A). Free abdominal somites with a median elevation in somites 3 to 6 (Figure 2 B).




Figure 2. *Teleophrys pococki* Rathbun, 1924. (A) Male in dorsal view; (B) ventral view; (C) right first gonopod in pleonal view, in stereomicroscopy (Scale bars = 5mm (A,B), 1mm (C).

The first gonopod (G1) is long and robust, base with plumose setae, lateral margin with simple setae sparsely distributed, a suture present from the base to the distal margin of the apical plate, rounded protuberance on the lateral margin; apical plate with two lobes covered with several acute spines (Figure 2C).

Originally, the species *T. pococki* was designated as a new species in a revision of the type material of *T. cristulipes* provided by Rathbun (1924), being these both closely related species, which can be easily separated by main characters: rostrum short and large, slightly bifid and not reaching the first antennal peduncle in *T. pococki* vs. rostrum with frontal margin large, bifid and overcoming the first antennal peduncle in *T. cristulipes*; carapace with only one antero-lateral tooth in *T. pococki* vs. carapace with 3-4 antero-lateral teeth in *T. cristulipes*; carapace smooth, with small granules and tubercles only in the branchial and cardiac regions in *T. pococki* vs. all carapace regions covered with small granules and tubercles in *T. cristulipes*. Ambulatory legs are slender, with tubercles forming a crest in the superior margin in *T. pococki* vs. ambulatory legs robust with a visible lobe at propodus in *T. cristulipes* (see more details in RATHBUN, 1925). Rathbun (1900), analyzing samples from Brazil, indicated *T. cristulipes* as occurring in the region; however, the same author posteriorly described the Brazilian species as *T. pococki*, indicating *T. cristulipes* as native from the Pacific Ocean. In addition, the distribution of *T. cristulipes* covers the Pacific Ocean, Mexico (Baja California), Costa Rica (Golfo Dulce), Panama, Colombia, Ecuador (Galapagos) (HOLTHUIS, 1979; HENDRICKX, 2010).

The species *T. pococki* is restricted to Western Atlantic, occurring in the United States (Louisiana), Bonaire, Curaçao, Honduras, Bahamas, Lesser Antilles, Colombia (Santa Marta), Venezuela and Brazil: (Amapá [this study], Rocas Atoll, Fernando de Noronha Archipelago, Pernambuco, Alagoas, Bahia and São Paulo, Trindade and Martin Vaz Archipelago) (GARTH, 1978; GOUVÊA, 1986; COELHO; ALMEIDA; BEZERRA, 2008; MELO, 1996; 1998; ALVES et al., 2012; FLEEGER; COWAN-JR.; PASCAL, 2013; CARMONA-SUÁREZ; POUPIN, 2016). In Brazilian waters, only two *Teleophrys* species (*T. pococki* and *T. ornatus*) have been reported; however, *T. ornatus* is recorded only in Fernando de Noronha Archipelago, Bahia and São Paulo (MELO, 1996; ALVES et al., 2012). The main differences between these two species are: *T. pococki* has the carapace unarmed, covered with granules and wider than long, and the ambulatory legs with a crest only on the upper surface; while *T. ornatus* has the carapace armed with tubercles and small spines and longer than wide, and the ambulatory legs showing a crest on both sides (superior and inferior margins) (see MELO, 1996, pages. 248, 249).

The mithracid species *T. pococki* is considered an ecosystem engineer, especially living in association with coral reefs (e.g. *Mussismilia hispida* (Verrill, 1901) in shallow habitats), or living in burrows in dead coral or artificial substrates such as artificial reefs, oil platforms and ship hulls (NOGUEIRA; NEVES; JOHNSON, 2015).

Furthermore, this species may be found in calcareous algae beds; nonetheless, it is commonly found using seaweed of the genera *Centroceras* Kützting, 1841, *Ceramium* Roth, 1797 and *Chaetomorpha* Kützting, 1845 as ornamentation on areas of the carapace and walking legs (GOUVÊA, 1986; LÓPEZ; SOLANO, 2005; TAVARES; MENDONÇA JR., 2022). Studies performed by Vera-Caripe et al. (2017) indicated the commensal association of *T. pococki* with the sea urchin *Tripneustes ventricosus* (Lamarck, 1816), being the echinoderm a protection against predators.

Into our grapsid specimen, we collected one male of *E. americanus* (Figure 3 A-D) (cw. 7.10 mm, cl. 7.50 mm and ww. 0.24 g) (Voucher Number: 22.2.1 A), at 85 m of depth, in areas of the GARS (September 2025), state of Amapá (Brazil) (03°00'27" N; 049°06'20" W), associated with rhodoliths in the gravel bottom. Our specimen agrees with the original description provided by A. Milne-Edwards (1880) and additional posterior descriptions of A. Milne-Edwards and Bouvier (1894), Rathbun (1918), Türkay (1975) and Melo (1996).

The main characteristics for the identification of this species are: carapace subquadrate, flattened posteriorly, with the surface covered with small granules and clumps of stout setae spread non-uniformly, including numerous short setae covering the carapace. The bilobed front reaches the middle of the first antennular article (Figure 3A). Three antero-lateral acute spines (excluding the orbital), being the second more pronounced/acute than the others (Figure 3A-C).

Chelipeds thin, covered by stout setae and small granules, slightly curved downward. Merus of the chelipeds contain five acute spines on the ventral-distal angle. Ambulatory legs covered with stout setae and small granules; merus of pereopods 2 to 5 (P2-P5) ending in a sharp distal tooth, merus of the second ambulatory leg (P2) containing three to four subdistal teeth (a higher and sharper) (Figure 3B). Dactyl of P2-P5 with five to six robust spines, being the tip trifurcated (similar to a hook). Abdominal somites one and two are reduced, and somites 4-6 are slightly fused. First gonopod (G1) robust, slightly curved towards the mesial line, with sparse setae in its mid-distal portion and dense pubescence at the tip (see Figure 3D) (modified from TÜRKAY, 1975; MELO, 1996).




Figure 3. *Euchirops americanus* A. Milne-Edwards, 1880. (A) Male in dorsal view; (B) inferior margin of the merus of the second pair of ambulatory legs (P2); (C) Carapace; (D) right first gonopod in pleonal view, in stereomicroscopy (Scale bars = 5mm (A), 1mm (D)).

In the Western Atlantic, the species most closely related to *E. americanus* is *E. antillensis*. Both species occur in the same habitats (sympatric species), however they are morphologically distinguished according to Türkay (1975, pag. 113, Fig. 16) and Alves-Júnior et al. (2016, pag. 3, Fig. 1) following the characters: the individuals of *E. americanus* have three to four subdistal teeth on the inferior margin of the merus of the second pair of ambulatory legs (P2), being the first or second (sometimes the last) larger and higher pitched than the others; while in *E. antillensis* all subdistal teeth on the inferior margin of the merus of P2 are of the same size. Additionally, Türkay (1975) indicated that the anterolateral teeth of *E. americanus* are less developed than in *E. antillensis*. Studies developed by Poupin (1994; 2018) indicated this character as valid for male and female specimens identification.

Until now, several specimens were erroneously assigned as *E. americanus*, especially in Indo-Pacific regions; however, after the revision provided by Türkay (1975) and additional observations of Manning and Holthuis (1981); these authors indicated that only two *Euchirograpsus* species (*E. americanus* and *E. antillensis*) are restricted to the Western Atlantic. The distribution of *E. americanus* covers: Canada (Bay of Fundy), USA (towards the Carolina's in Toms/ Meys Canyon and Baltimore Canyon, Massachusetts in Canyon at the edge of Georges Bank, Delaware, New Jersey, Louisiana, Florida), Gulf of Mexico, Dry Tortugas, Cuba, Lesser Antilles, West Indies, Guadalupe, Barbados, Colombia, Venezuela, French Guiana and Brazil (Amapá [this study], Maranhão, Rio Grande do Norte and Rio Grande do Sul states) (ZARIQUIEY ALVAREZ, 1968; TÜRKAY, 1975; MELO, 1996; 1998; CORBARI et al. 2015; ALVES-JÚNIOR et al., 2016; WILSON; POHLE, 2016; POUPIN, 2024). According to Wilson and Pohle (2016), this species is normally found in tropical areas; however, possibly due to climate change events, *E. americanus* has expanded into subtropical areas, as observed in cold waters (-8.3 °C) of Canada (Bay of Fundy).

The presence of these species in the State of Amapá may be associated with the GARS, once this region encompasses a wide range of micro- and macro habitats beneath the Amazon River plume, characterized by high energy flow and acting as connection between South America and the Caribbean Sea (MOURA et al., 2016; VALE et al., 2022; ALVES-JÚNIOR; MARTINS; CINTRA, 2024b). Along the GARS, the migration/distribution of invertebrate and vertebrate species may occur in a stepping-stones process, especially in areas below 70 m (ALVES-JÚNIOR; MARTINS; CINTRA, 2024b; MARTINS; ALVES-JÚNIOR; CINTRA, 2025). This explains the widespread occurrence of *T. pococki* and *E. americanus* throughout the Western Atlantic.

Two other additional hypotheses that may explain the presence of both species in northern Brazil are: 1) their transport (larvae and adults) is associated with ships on route between Brazil and the Caribbean Sea; 2) these species were probably already distributed throughout the Amazon continental shelf region, but the lack of researchers and specialized sampling efforts along the GARS have hindered knowledge of the distribution of *T. pococki* and *E. americanus*.

4 Conclusions

Despite the high biological diversity found in the ACS, the region is suffering severe anthropogenic impacts, especially associated with fishing activities, climate change, marine debris and ghost fishing. As a result, several species of marine invertebrates and vertebrates are included in threatened species lists, even before they have been catalogued or their biological characteristics fully understood. The presence of only a single specimen of *T. pococki* and *E. americanus* in the northern region limits our understanding of their biology and conservation status. In conclusion, we herein update the geographic distribution of two brachyuran crab species from the Amazon continental shelf; however, further studies in the Amazon reef region are needed to improve the knowledge of crustacean diversity in northern Brazil.

CREDIT AUTHORSHIP CONTRIBUTION STATEMENT

Conceptualization: D.E.G.M., I.H.A.C., F.A.A.J. and B.B. Funding: I.H.A.C., F.A.A.J. and B.B. Data curation, Methodology: D.E.G.M., I.H.A.C. and F.A.A.J. Investigation: D.E.G.M. and F.A.A.J. Visualization: D.E.G.M. Writing - original draft: D.E.G.M., I.H.A.C. and F.A.A.J. Writing - Review & Editing: D.E.G.M. and B.B.

DECLARATION OF INTEREST

The authors disclose that they have no known competing financial interests or personal relationships that could have appeared to influence the study reported in this manuscript.

FUNDING SOURCE

This research was financed by the *Conselho Nacional de Desenvolvimento Científico e Tecnológico* - CNPq (Brazil), through the productivity grant award, reference numbers: 304401/2025-0 (author Israel Cintra) and 302175/2025-3 (Bianca Bentes).

ETHICAL PROCEDURES

No live animal treatments were carried out in this study. All specimens were retrieved from regulated commercial fishing operations and were dead upon examination.

ACKNOWLEDGEMENTS

The authors would like to thank the *Centro Nacional de Pesquisa e Conservação da Biodiversidade Marinha do Norte* (CEPNOR, Brazil) for the sampling of the examined material and the support in the laboratory. Additionally, we would like to thank the anonymous reviewers for their valuable comments throughout the manuscript.

REFERENCES

- ALVES, D.F.; BARROS-ALVES, S.P.; TEIXEIRA, G.M.; COBO, V.J. Mithracinae (Decapoda: Brachyura) from the Brazilian coast: Review of the geographical distribution and comments on the biogeography of the group. *Nauplius*, v. 20, n. 1, p. 51-62, 2012. Available from: <https://www.scielo.br/j/nau/a/ZxwV8fKNxv65FMVvKRxNH7J/?format=pdf&lang=en>. Accessed on: 15 Sep 2025.
- ALVES-JÚNIOR, F.A.; MARTINS, D.E.G.; KLAUTAU, A.G.C.M.; BARROS, A.P.C.; CINTRA, I.H.A. Pelagic *Sargassum* spp. blooms as an agent of marine litter transport along the Amazon continental shelf. *Acta Fisheries and Aquatic Resources*, v. 12, n. 2, p. 121-126, 2024a. Available from: <https://doi.org/10.46732/actafish.2024.12.2.121-126>
- ALVES-JÚNIOR, F.A.; MARTINS, D.E.G.; CINTRA, I.H.A. Becoming a mother too early: The first record of the smallest ovigerous female of the deep-sea squat lobster *Iridonida iris* (A. Milne-Edwards, 1880) (Decapoda: Munidiidae). *Pan-American Journal of Aquatic Science*, v. 19, n. 2, p. 92-95, 2024b. Available from: <https://doi.org/10.54451/PanamJAS.19.2.92>
- ALVES-JÚNIOR, F.A.; SANTANA, J.L.; ARAÚJO, M.S.L.C.; SOUZA-FILHO, J.F. First record of *Euchirograpsus americanus* A. Milne-Edwards, 1880 (Crustacea: Decapoda: Plagusiidae) from northeastern Brazil. *Nauplius*, v. 24, p. e2016023, 2016. Available from: <https://doi.org/10.1590/2358-2936e2016023>
- BESCHIN, C.; BUSULINI, A.; TESSIER, G.; ZORZIN, R. I crostacei associati a coralli nell'Eocene inferiore dell'area di Bolca (Verona e Vicenza, Italia nordorientale). *Memorie del Museo Civico di Storia Naturale di Verona, serie 2, Sezione Scienze della Terra* 9, 189 p., 2016. Available from: https://www.valdalpnoneheritage.it/ATS_Documenti/2016_I%20Crostacei%20associati%20a%20coralli.pdf. Accessed on: 20 Oct 2025.
- CANO, G. Viaggio della R. Corvetta Vettor Pisani attorno al globo. - Crostacei Brachiuri ed Anomuri. *Bollettino della Società di naturalisti in Napoli*, v. 3, p. 169-268, 1889. Available from: <https://www.marinespecies.org/aphia.php?p=sourcedetails&id=391827>. Accessed on: 15 Oct. 2025.
- CARMONA-SUÁREZ, C.; POUPIN, J. Majoidea crabs from Guadeloupe Island, with a documented list of species for the Lesser Antilles (Crustacea, Decapoda, Brachyura, Majoidea). *Zoosystema*, v. 38, n. 3, p. 353-387, 2016. Available from: <https://doi.org/10.5252/z2016n3a5>
- CASTAÑO, N.C.; CAMPOS, N.H. Los cangrejos Araña (Decapoda: Brachyura: Majoidea) del Caribe colombiano. *Biota Colombiana*, v. 4, n. 2, p. 261-269, 2003. Available from: <https://www.redalyc.org/articulo.oa?id=49140208>. Accessed on: 17 Sep 2025.
- COELHO, P.A.; ALMEIDA, A.O.; BEZERRA, L.E.A. Checklist of the marine and estuarine Brachyura (Crustacea: Decapoda) of the northern and northeastern Brazil. *Zootaxa*, v. 1956, n.1, p. 1-58, 2008. Available from: <https://doi.org/10.11646/zootaxa.1956.1.1>
- CORBARI, L.; MARTIN-LEFÈVRE, P.; SABROUX, R.; SOUBZMAIGNE, S. Crustacés. In: PASCAL, O.; TOUROULT, J.; BOUCHET, P. (Eds.). *Expédition La Planète Revisitée Guyane 2014-15: Synthèse des Premiers Résultats*. Muséum national d'Histoire naturelle. p. 162-174, 2015. Available from <https://inpn.mnhn.fr/docs-web/docs/download/249749>. Accessed on: 17 Sep 2025.
- DANA, J.D. Crustacea Grapsoidea, (Cyclometopa, Edwardsii): Conspectus Crustacearum quæ in Orbis Terrarum circumnavigatione, Carolo Wilkes e classe Reipublicæ Fœderatæ Duce, lexit et descriptis J. D. Dana. *Proceedings of the Academy of Natural Sciences of Philadelphia*, v. 5, p. 247-254. Available from: <https://www.biodiversitylibrary.org/part/22775>. Accessed on: 05 Nov 2025.
- DecaNet (World List of Decapoda). DecaNet. Mithracidae MacLeay, 1838. 2025a. Available from: <https://www.decanet.info/aphia.php?p=taxdetails&id=439056>. Accessed on: 12 Nov 2025.
- DecaNet (World List of Decapoda). DecaNet. Plagusiidae Dana, 1851. 2025b. Available from: <https://www.decanet.info/aphia.php?p=taxdetails&id=439056>. Accessed on: 12 Nov 2025.
- DRAKE, T.W.; HEMINGWAY, J.D.; KUREK, M.R.; PEUCKER-EHRENBRINK, B.; BROWN, K.A.; HOLMES, R.M.; VALIER, G.; MOURA, J.M.S.; MITSUYA, M.; WASSENAAR, L.I.; SIX, J.; SPENCER, R.G.M. The pulse of the Amazon: Fluxes of dissolved organic carbon, nutrients, and ions from the world's largest river. *Global Biogeochemical Cycles*, v.

35, n. 4, p. e2020GB006895, 2021. Available from: <https://doi.org/10.1029/2020GB006895>

FLEEGER, S.T.D.J.; COWAN-JR., J.H.; PASCAL, P.-Y. What Is the Relative Importance of Phytoplankton and Attached Macroalgae and Epiphytes to Food Webs on Offshore Oil Platforms? *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, v. 5, p. 53-64, 2013. Available from: <https://doi.org/10.1080/19425120.2013.774301>

FUJITA, Y.; NARUSE, T. A new genus and species of a submarine cave crab of the family Plagusidae Dana, 1851 (Crustacea: Brachyura: Grapsoidea) from Okinawa Island, Ryukyu Islands, southwestern Japan. *Zootaxa*, v. 5410, n. 3, p. 408-418, 2024. Available from: <https://doi.org/10.11646/zootaxa.5410.3.8>

GARTH, J.S. Marine Biological Investigations in the Bahamas 19. Decapoda Brachyura. *Sarsia*, v. 63, n. 4, p. 317-333, 1978. Available from: <https://doi.org/10.1080/00364827.1978.10411352>

GOUVÊA, E.P. A carcinofauna do litoral rochoso de Salvador, BA, e alguns aspectos ecológicos. *Ciência e Cultura*, v. 38, n. 2, p. 346-354, 1986. Available from: <http://memoria.bn.gov.br/docreader/003069/45802>. Accessed on: 08 Oct 2025.

GUINOT, D. A new genus of the family Plagusidae Dana, 1851, close to *Plagusia* Latreille, 1804 (Crustacea, Decapoda, Brachyura). *Zootaxa*, v. 1498, n. 1, p. 27-33, 2007. Available from: <https://doi.org/10.11646/zootaxa.1498.1.2>

HENDRICKX, M.E. On a small collection of decapod crustaceans (Malacostraca, Decapoda) from the Galapagos Islands, collected by P. H. Vercammen-Grand-Jean, in 1964. *Crustaceana*, v. 83, n. 8, p. 1009-1015, 2010. Available from: <https://doi.org/10.1163/001121610X502902>

HOLTHUIS, L.B. A small collection of Decapod Crustacea from Galapagos Islands. *Spedizione L. Mares-G.R. T. S.*, Florence, Italy, 1979. Available from: <https://scispace.com/pdf/a-small-collection-of-decapod-crustacea-598sdro3bn.pdf>. Accessed on: 13 Nov 2025.

KLAUTAU, A.G.C.M.; CORDEIRO, A.P.B.; CHAGAS, R.A.; SANTOS, W.C.R.; MARCENIUK, A.P.; NÓBREGA, P.S.V.; MARTINELLI-LEMOES, J.M.; CINTRA, I.H.A.; CASTRO, N.S.S.; BASTOS, C.E.M.C.; ALVES-JÚNIOR, F.A.; PINHEIRO, L.C.; BENTES, B. Biodiversity hotspots and threatened species under human influence in the Amazon continental shelf. *Scientific Reports*, v. 15, p. 26681, 2025. Available from: <https://doi.org/10.1038/s41598-025-11261-x>

KÜTZING, F.T. (1842 '1841'). Ueber *Ceramium* Ag. *Linnaea*, v. 15, n. 6, p. 727-746, 1841. Available from: https://www.algaebase.org/search/bibliography/detail/?biblio_id=11288. Accessed on: 15 Nov 2025.

KÜTZING, F.T. *Phycologia Germanica*, d.i. Deutschlands Algen in bündigen Beschreibungen. Nebst einer Anleitung zum Untersuchen und Bestimmen dieser Gewächse für Anfänger. Nordhausen, 1845, 340 pp., Available from: <https://doi.org/10.5962/bhl.title.13687>

LAMARCK, J.B.M. *Histoire naturelle des animaux sans vertèbres*, Tome troisième, Paris, Deterville/Verdière, 586 p., 1816. Available from: <https://doi.org/10.5962/bhl.title.12712>

LATREILLE, P.A. Tableau méthodique des crustacés. Classe septième. Crustacés (1), Crustacea. *Nouveau Dictionnaire d'Histoire Naturelle, Appliquée aux Arts, Principalement à l'Agriculture et à l'Economie Rurale et Domestique: par une Société de Naturalistes et d'Agriculteurs: avec des figures tirées des trois Règnes de la Nature*. v. 24, p. 123-127, 1804. Available from: <https://doi.org/10.5962/bhl.title.151939>

LÓPEZ, M.M.; SOLANO, O.D. Estado poblacional de *Echinometra lucunter* (Echinoida: Echinometridae) y su fauna acompañante en el litoral rocoso del Caribe Colombiano. *Revista de Biología Tropical*, v. 53, n. 3, p. 291-297, 2005. Available from: <https://tropicalstudies.org/rbt/attachments/suppls/sup53-3%20echinoderm/19-MONROY-Asp.pdf>. Accessed on: 05 Nov 2025.

MACLEAY, W.S. Illustrations of the Annulosa of South Africa. On the Brachyurous Decapod Crustacea. Brought from the Cape by Dr. Smith. In: Smith, A. (Ed.), *Illustrations of the zoology of South Africa: consisting chiefly of figures and descriptions of the objects of natural history collected during an expedition into the interior of South Africa, in the years 1834, 1835, and 1836, fitted out by "The Cape of Good Hope Association for Exploring Central Africa": together with a summary of African zoology, and an inquiry into the geographical ranges of species in that quarter of the globe*. London: Authority of the Lords Commissioners of Her Majesty's Treasury, 1838. Vol. 5. p. 53-71. Available from: <https://www.biodiversitylibrary.org/item/41545>. Accessed on: 05 Nov 2025.

MANNING, R.B.; HOLTHUIS, L.B. West African brachyuran crabs (Crustacea: Decapoda). *Smithsonian Contributions to Zoology*, v. 306, p. 1-379, 1981. Available from: <https://doi.org/10.5479/si.00810282.306>

MARTINS, D.E.G.; ALVES-JÚNIOR, F.A.; CINTRA, I.H.A. New record of the swimming crab *Portunus sayi* (Gibbes, 1850) (Decapoda: Portunidae) from the Amazon

continental shelf. *Environmental Smoke*, v. 8, p. es20250003, 2025. Available from: <https://doi.org/10.32435/envsmoke-2025-0003>

MELO, G.A.S. *Manual de identificação dos Brachyura (caranguejos e siris) do litoral brasileiro*. Editora Plêiade, São Paulo, 603 pp, 1996. Available from: <https://repositorio.usp.br/item/000921427>. Accessed on: 10 Oct. 2025.

MELO, G.A.S. Malacostraca-Eucarida. Brachyura. Oxyrhyncha and Brachyrhyncha. p. 455-515. In: YOUNG, P.S. (ed). *Catalogue of Crustacea of Brazil*. Rio de Janeiro, Museu Nacional, Série Livros 6, 1998. Available from: <https://www.scielo.sa.cr/scieloOrg/php/reflinks.php?refpid=S0034-7744200800030002100053&lng=es&pid=S0034-77442008000300021>. Accessed on: 18 Oct. 2025.

MILNE-EDWARDS, A. Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico and in the Caribbean Sea, 1877, 1878, 1879, by the U.S. Coast Survey Steamer "Blake", Lieut-Commander C.D. Sigsbee, U.S.N., and Commander J.R. Bartlett, U.S.N., Commanding. VIII. Études préliminaires sur les Crustacés, 1re Partie. *Bulletin of the Museum of Comparative Zoology at Harvard College*, v. 8, n. 1, p. 1-68, 1880. Available from: <https://biostor.org/reference/106168>. Accessed on: 05 Nov 2025.

MILNE-EDWARDS, A.; BOUVIER, E.L. Crustacés décapodes provenant des campagnes du yacht l'Hirondelle (1886, 1887 et 1888). Première partie: Brachyures et Anomoures. *Résultats des Campagnes Scientifique Monaco*, v. 7, p. 1-112, 1894. Available from: <https://www.biodiversitylibrary.org/item/31107>. Accessed on: 15 Oct 2025.

MILNE-EDWARDS, H. Mémoire sur la famille des ocyropodiens. *Annales des Sciences Naturelles*, v. 20, p. 163-228, 1853. Available from: <https://biostor.org/reference/160231>. Accessed on: 16 Oct 2025.

MOURA, R.L.; AMADO-FILHO, G.M.; MORAES, F.C.; BRASILEIRO, P.S.; SALOMON, P.S.; MARQUES, M.M.; BASTOS, A.C.; ALMEIDA, M.G.; SILVA-JÚNIOR, J.M.; ARAÚJO, B.F.; BRITO, F.P.; RANGEL, T.P.; OLIVEIRA, B.C.V.; BAHIA, R.G.; PARANHOS, R.P.; DIAS, R.J.S.; SIEGLE, E.; FIGUEIREDO-JÚNIOR, A.G.; PEREIRA, R.C.; LEAL, C.V.; HAJDU, E.; ASP, N.E.; GREGORACCI, G.B.; NEUMANN-LEITÃO, S.; YAGER, P.L.; FRANCINI-FILHO, R.B.; FROÉS, A.; CAMPEÃO, M.; SILVA, B.S.; MOREIRA, A.P.B.; OLIVEIRA, L.; SOARES, A.C.; ARAÚJO, L.; OLIVEIRA, N.L.; TEIXEIRA, J.B.; VALLE, R.A.B.; THOMPSON, C.C.; REZENDE, C.E.; THOMPSON, F.L. An extensive reef system at the Amazon River mouth.

Science Advances, v. 2, n. 4, p. 1-11, 2016. Available from: <https://doi.org/10.1126/sciadv.1501252>

NG, P.K.L.; GUINOT, D.; DAVIE, P.J.F. *Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world*. *The Raffles Bulletin of Zoology*, v. 17, n. 1, p. 1-286, 2008. Available from: <https://lcnhm.nus.edu.sg/wp-content/uploads/sites/11/app/uploads/2017/04/s17rbz.pdf>. Accessed on: 03 Nov 2025.

NOGUEIRA, M.M.; NEVES, E.; JOHNSON, R. Effects of habitat structure on the epifaunal community in *Mussismilia* corals: does coral morphology influence the richness and abundance of associated crustacean fauna?. *Helgoland Marine Research*, v. 69, p. 221-229, 2015. Available from: <https://doi.org/10.1007/s10152-015-0431-x>

PORTELL, R.W.; COLLINS, J.S.H. Decapod crustaceans of the Lower Miocene Montpellier Formation, White Limestone Group of Jamaica. *Cainozoic Research*. v. 3, n. 1-2, p. 109-126, 2004. Available from: <https://natuurtijdschriften.nl/pub/541705/CR2004003001007.pdf>. Accessed on: 15 Oct 2025.

POUPIN, J. *Faune marine profonde des Antilles françaises: récoltes du navire Polka faites en 1993*. Paris: ORSTOM, 1994. Collection Études et Thèses, 1994. Available from: https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_2/etudes_theses/39603.pdf. Accessed on: 16 Sep 2025.

POUPIN, J. *Les Crustacés décapodes des Petites Antilles*. Avec de nouvelles Observations pour Saint-Martin, la Guadeloupe et la Martinique. Muséum National d'Histoire Naturelle, Patrimoines naturels 77, 2018. Available from: <https://sciencepress.mnhn.fr/sites/default/files/documents/en/cpn77-listedoc.pdf>. Accessed on: 12 Nov 2025.

POUPIN, J. A documented checklist of the Crustacea (Stomatopoda, Decapoda) of the southern Guianas (Guyana, Suriname, French Guiana, Brazil Amapá). *European Journal of Taxonomy*, v. 954, p. 1-197, 2024. Available from: <https://doi.org/10.5852/ejt.2024.954.2653>

RATHBUN, M.J. The decapod and stomatopod Crustacea. I. Results of the Branner-Agassiz Expedition to Brazil. *Proceedings of the Washington Academy of Sciences*, v. 2, p. 133-156, 1900. Available from: <https://www.biodiversitylibrary.org/page/8872151#page/172/mode/1up>. Accessed on: 06 Nov 2025.

RATHBUN, M.J. *Investigations of the Aquatic Resources and Fisheries of Porto Rico by the United States Fish*

Commission Steamer Fish Hawk in 1899. The Brachyura and Macrura of Porto Rico. **Bulletin of the United States Fish Commission**, v. 20, p. 1-127, 1901. Available from: <https://www.decanet.info/aphia.php?p=sourcedetails&iid=125904>. Accessed on: 12 Nov 2025.

RATHBUN, M.J. The grapsoid crabs of America. **United States National Museum Bulletin**, v. 97, p. 1-461, 1918. Available from: <https://repository.si.edu/items/239cd4de-44e5-424d-93b9-3d72ed7813dd>. Accessed on: 8 Nov 2025.

RATHBUN, M.J. New species and subspecies of spider crabs. **Proceedings of the United States National Museum**, v. 64, n. 2504, p. 1-5, 1924. Available from: <https://repository.si.edu/server/api/core/bitstreams/5a41306b-fbab-40f8-bb32-2db44a44f9f7/content>. Accessed on: 03 Nov 2025.

RATHBUN, M.J. The spider crabs of America. **United States National Museum Bulletin**, v. 129, p. 1-613, 1925. Available from: <https://repository.si.edu/items/d54c874d-86aa-4a49-8481-188e9442518b>. Accessed on: 06 Nov 2025.

ROTH, A.W. *Catalecta Botanica* (Quibus plantae novae et minus cognitae describuntur atque illustrantur). Fasc. 1. Lipseae., 244 p., 1797. Available from: <https://www.biodiversitylibrary.org/page/56933819>. Accessed on: 11 Nov 2025.

SCHUBART, C.D.; CUESTA, J.A. Phylogenetic relationships of the Plagusidae Dana, 1851 (Brachyura), with description of a new genus and recognition of Percnidae Stevcic, 2005, as an independent family. In: CASTRO, P.; DAVIE, P.J.F.; NG, P.K.L.; FORGES, B.R. **Studies on Brachyura: a Homage to Danièle Guinot**. Crustaceana Monographs 11. Leiden: Brill. p. 279-300, 2010. Available from: <https://decapoda.nhm.org/pdfs/31482/31482.pdf>. Accessed on: 12 Oct 2025.

STIMPSON, W. Sketch of a revision of the genera of Mithracidae. **The American Journal of Science and Arts**, Series 2, v. 29, n. 85, p. 132-133, 1860. Available from: <https://archive.org/details/sketchofrevision00stim/page/132/mode/2up>. Accessed on: 28 Oct 2025.

STIMPSON, W. Preliminary report on the Crustacea dredged in the Gulf Stream in the Straits of Florida, by L. F. de Pourtales, Assist. U. S. Coast Survey. Part I. Brachyura. **Bulletin of the Museum of Comparative Zoölogy**, v. 2, n. 2, p. 109-160, 1871. Available from: <https://www.biodiversitylibrary.org/page/6313618#page/125/mode/1up>. Accessed on: 18 Oct 2025.

TAVARES, M.; MENDONÇA Jr., J.B. Brachyuran crabs (Crustacea, Decapoda) from the remote oceanic Archipelago Trindade and Martin Vaz, South Atlantic Ocean. **Zootaxa**, v. 5146, n. 1, p. 1-129, 2022. Available from: <https://doi.org/10.11646/zootaxa.5146.1.1>

TÜRKAY, M. Zur Kenntniss der Gattung Euchiropsus mit Bemerkungen zu Brachyropsus und Litocheira (Crustacea, Decapoda). **Senckenbergiana biologica**, v. 56, n. 1, p. 103-132, 1975. Available from: <https://decapoda.nhm.org/pdfs/12714/12714.pdf>. Accessed on: 14 Oct 2025.

TÜRKAY, M. Zwei neue Grapsiden-Arten aus dem Indopazifik (*Euchiropsus madagascariensis* n. sp. und *Miersiropsus australiensis* n. sp.) mit Einführung von *Miersiropsus* n. gen. (Crustacea: Decapoda: Grapsidae). **Senckenbergiana biologica**, v. 59, n. 1-2, p. 133-141, 1978. Available from: <https://decapoda.nhm.org/pdfs/23688/23688.pdf>. Accessed on: 11 Oct 2025.

VALE, N.F.; BRAGA, J.C.; MOURA, R.L.; SALGADO, L.T.; MORAES, F.C.; KAREZ, C.S.; CARVALHO, R.T.; SALOMON, P.S.; MENANDRO, P.S.; AMADO-FILHO, G.M.; BASTOS, A.C. Distribution, morphology and composition of mesophotic 'reefs' on the Amazon continental margin. **Marine Geology**, v. 447, p. 106779, 2022. Available from: <https://doi.org/10.1016/j.margeo.2022.106779>

VERA-CARIPE, J.; DIAZ, O.; LIRA, C.; BOLAÑOS, J. Crustáceos decápodos asociados a *Tripterygion* (*Tripterygion*) (*Echinodermata*; *Echinoidea*) de la isla La Borracha, parque nacional Mochima, Venezuela. **Boletín del Instituto Oceanográfico de Venezuela**, v. 56, n. 1, p. 61-68, 2017. Available from: <http://saber.udo.edu.ve/index.php/boletiniov/article/view/2644/1566>. Accessed on: 04 Nov 2025.

VERRILL, A. E. Variations and nomenclature of Bermudian, West Indian and Brazilian reef corals, with notes on various Indo-Pacific corals. **Transactions of the Connecticut Academy of Arts and Sciences**, v. 11, p. 63-168, 1901. Available from: <https://www.biodiversitylibrary.org/page/13464046>. Accessed on: 25 Oct 2025.

WEBER, S.C.; CARPENTER, E.J.; COLES, V.J.; YAGER, P.L.; GOES, J.; MONTOYA, J.P. Amazon River influence on nitrogen fixation and export production in the western tropical North Atlantic. **Journal of Oceanology and Limnology**, v. 62, p. 618-631, 2017. Available from: <https://doi.org/10.1002/lno.10448>

WILSON, B.M.; POHLE, G.W. Northern range expansion of the American talon crab, *Euchirograpsus americanus* A. Milne-Edwards, 1880 (Decapoda, Grapsoidea, Plagusiididae), to the Bay of Fundy, Canada. **Crustaceana**, v. 89, n. 2, p. 163-173, 2016. Available from: <https://www.jstor.org/stable/43965981>. Accessed on: 18 Oct 2025.

WINDSOR, A.M.; FELDER, D.L. Molecular phylogenetics and taxonomic reanalysis of the family Mithracidae MacLeay (Decapoda: Brachyura: Majoidea). **Invertebrate Systematics**, v. 28, p. 145-173, 2014. Available from: <https://doi.org/10.1071/IS13011>

ZARIQUIEY-ALVAREZ, R. Crustáceos decápodos ibéricos. **Investigación Pesquera, Barcelona**, v. 32, p. 1-510, 1968. Available from: <https://decapoda.nhm.org/pdfs/10992/10992.pdf>. Accessed on: 22 Oct 2025.