



PRESENCE OF METACERCARIAE OF *Opisthorchis* sp. BLANCHARD, 1895 (TREMATODA: OPISTHORCHIIDAE) IN NEW HOST *Macrobrachium amazonicum* (HELLER, 1862), FROM THE NORTHERN BRAZIL: AN EMERGING RISK TO HUMANS BY THE SHRIMP CONSUMPTION

Déborah Elena Galvão Martins^{1,2*}; Israel Hidenburgo Aniceto Cintra²; Mayara Galvão Martins³; Flavio de Almeida Alves-Júnior^{1,2}

Abstract

Herein, we report the first observation of the metacercaria *Opisthorchis* sp. Blanchard, 1895 recovered from a new intermediate host *Macrobrachium amazonicum* (Heller, 1862), an important food resource from northern Brazil, emphasising the risk of contamination from inappropriate human consumption. The shrimp specimens of *M. amazonicum* were collected during June 2023, using a local trap called matapi in the Guamá River ($01^{\circ}27'54.2''S$; $048^{\circ}26'02.6''W$), which supplies the city of Belém, showing a characteristic of an eutrophicated river. The collected specimens were sexed, measured, weighted and dissected in the carapace region, to observe the presence of parasites in gills, hepatopancreas and gonads. We observed 56 specimens of *M. amazonicum* (20 males, 32 females and 4 ovigerous females), of which thirteen individuals (23.21% of total sampled) showed encysted metacercariae of *Opisthorchis* sp. only in gonadal tissues, with high preference in females (66.6%), when compared with males (33.4%). The observation of metacercaria in Amazon river prawn is a risk to human health, due to the parasite's life cycle being completed in humans, leading to several health problems such as hepatic and biliary degradation, diarrhoea, vomiting and respiratory problems. Additionally, the eutrophication process observed in the Guamá River may favour parasite proliferation, and dissemination to other hosts and regions, possibly increasing the cases of people infected with parasites in Belém metropolitan area. Based on that, this work is an alert to the authorities for the risk of contamination of the population due to inadequate consumption of (molluscs, crustaceans and fishes) from the eutrophic waters of the Guamá River and adjacent waters.

Keywords: Trematode parasite. New host association. Risk to human health. Fish-borne trematodes.

¹Universidade Federal do Pará, Núcleo de Ecologia Aquática e Pesca (NEAP), Rua Augusto Corrêa, s/n, Guamá, CEP: 66075-110, Belém, Pará, Brasil. Programa de Pós-Graduação em Ecologia Aquática e Pesca (PPGEAP) da Universidade Federal do Pará (UFPA), Belém, Pará, Brasil.

²Laboratório de Crustáceos (LABCRUS), Instituto Socioambiental e dos Recursos Hídricos (ISARH), Universidade Federal Rural da Amazônia (UFRA), Avenida Presidente Tancredo Neves, nº 2501, Terra Firme, CEP: 66077-830, Belém, Pará, Brasil.

³Instituto de Desenvolvimento Sustentável Mamirauá, Grupo de Pesquisa Inovação, Desenvolvimento e Adaptação de Tecnologias Sustentáveis, Estrada do Bexiga, nº 2584, CEP: 69553-225, Tefé, Amazonas, Brasil.

*Corresponding author:deborah.martins@ufra.edu.br

Submitted on: 17Jul. 2023
Accepted on: 5 Aug. 2023
Published on: 31 Aug. 2023

1 Introduction

The biodiversity in aquatic environments is correlated to abiotic (e.g. temperature, salinity, pH and dissolved oxygen); and biotic factors (e.g input of organic matter, photosynthesis (primary production) and energy flow in food chain), with the species playing a wide range of biological interactions such as: symbiosis, mutualism, competition, predation, commensalism and parasitism; this latter can be associated with the health of the environment and indications of environmental impacts (EMPARANZA et al., 2011; GÓNGORA-GÓMEZ et al., 2015).

The transmission of diseases through the ingestion of contaminated water or food is a public health problem in underdeveloped and developing countries, including transmission of parasites between aquatic animals and humans (BELIZARIO et al., 2001; HOP et al., 2007; PUMIDONMING et al., 2018).

The fish-borne trematodes (FBT) are one of the main parasitic diseases transmitted between aquatic organisms (molluscs and fishes) and birds, domestic animals and humans (SITHITHAWORN; HASWELL-ELKINS, 2003; LOVIS et al., 2009; CHAI et al., 2017).

Currently in literature, the trematodes use snails as first intermediate host (metacercaria stage), freshwater fishes as second intermediate host; these are species consumed by fish-eating birds and mammals (including humans), which become definitive hosts (TOUCH et al., 2009; DAO et al., 2017; PHYO MYINT et al., 2020).

Up to date, more than 20 million people are contaminated with trematodes, especially in Asian and Middle Eastern countries, where the degradation of water bodies has been intensified in the last years, causing an increase in the number of cases by the consumption of raw fish and molluscs (PETNEY et al., 2013; PHYO MYINT et al., 2020).

The trematode family Opisthorchiidae Looss, 1899 is composed of 53 genera distributed worldwide, occurring as obligate parasites in a wide range of hosts. One of the most diverse genus into Opisthorchiidae is *Opisthorchis* Blanchard, 1895, covering 47 species (CHAI, 2007; JEON et al., 2012; NOUDEH; PAZOOKI, 2019; WORMS, 2023).

Its proliferation around the world is caused by the inadequate human consumption of aquatic organisms; the parasite can reach great distances inhabiting the human intestines, additionally observed in migratory birds, fishes and by the aquarium trade (LEMOS et al., 2007).

In Brazil, the presence of these trematodes can be associated with water quality, the tourism flows in parallel with the lack of basic sanitation in some regions (THATCHER, 1993; PINTO; MELO, 2012; MATTOS; VERICIMO; SÃO CLEMENTE, 2013); aquarium trade process and the transport of these parasites through the introduction of exotic species, as observed by Pinto; Melo (2010) through the introduction of snail *Melanoides tuberculata* (Müller, 1774), an intermediate host of trematodes *Centrocestus formosanus* (Nishigori, 1924).

Despite being widely reported in the world, in Brazil, studies with trematodes are poorly documented, especially in the northern region; crustaceans are usually studied for growth, reproduction and fishery parameters, but not studied as parasite hosts. This study is the first observation of metacercariae of *Opisthorchis* sp. Blanchard, 1895 recovered from a new intermediate host *Macrobrachium amazonicum* (Heller, 1862), an important food resource from northern Brazil, emphasising the risk of contamination from inappropriate human consumption.

2 Material and Methods

Study Area

The prawn specimens of *M. amazonicum* were collected in the Guamá River ($01^{\circ}27'54.2''S$; $048^{\circ}26'02.6''W$) (Figure 1), one of the main rivers that crosses the city of Belém, in the State of Pará. The river is highly eutrophic, with muddy and turbid waters, with a strong presence of clay particles in suspension, receiving domestic waste from the Belém metropolitan area, as well as chemical elements from the factories located along the Guamá River (COSTA et al., 2022). This region presents a tropical humid climate (Köppen classification Af), with high temperatures all over the year (always above $18^{\circ}C$) and annual precipitation between 2250 and 3000 mm, with the characteristic rainy period between December and May and the dry period between June and November. The samples occurred in June 2023, using a local trap called "matapi" placed at night time and recovered in the next morning.

Laboratory Procedures

The collected individuals were transported alive to the carcinology laboratory (LABCRUS) of the Federal Rural University of the Amazon (UFRA), where the individuals were identified, sexed and measured with a calliper (0.01 mm) in Total Length (TL), Carapace Length (CL) and Total Weight (TW). Afterwards, the specimens were dissected in the carapace region for parasites in gills, hepatopancreas and gonads under a stereomicroscope.




Figure 1. Map of the study area indicating the urbanised area, corresponding to the metropolitan region of Belém- PA. Black star = sampling point.

In specimens that showed trematodes, the tissues were removed, transferred to petri dishes containing saline solution (NaCl 0.85%) and then analysed by means of optical microscopy to determine whether metacercariae were present for counting and photographing using the camera Motic Moticam5. For the fixation of metacercariae, we used AFA (Acetone - Formaldehyde - Alcohol).

For the identification of metacercariae, we followed the descriptions provided by Vajrasthira; Harinasuta; komiya (1961) and Scholz; Ditrich; Giboda (1992), for individuals of the genus: cyst in elliptical form, with equal size of oral sucker and ventral sucker, brownish pigment granules scattered within the body and an O-shaped excretory bladder occupying the greater part of the posterior body.

3 Results and Discussion

We collected 56 specimens of *M. amazonicum* (Figure 2A), being 20 males, 32 females and 4 ovigerous females, measuring: TL (min.: 6.18; max.: 9.85 cm; mean: 6.18 ± 1.12), CL (min.: 0.92; max.: 2.44 cm; mean: 1.45 ± 0.326) and TW (min.: 0.34; max.: 6.83 g; mean: 1.80 ± 1.04), which 13 individuals (23.21% of total sampled) showed the metacercariae of *Opisthorchis* sp. between the gonadal tissue (Figure 2B, C, D).

There were 53 metacercariae (covering between 3 and 5 metacercariae by prawn), with these parasites showing sizes between 288 and 725 μm (mean: $400 \mu\text{m} \pm 121$). The highest preference of *Opisthorchis* sp. was observed in female specimens of *M. amazonicum* (66.6%), when compared with males (33.4%).




Figure 2. (A) *Macrobrachium amazonicum* (Heller, 1862) parasitised by metacercariae of *Opisthorchis* sp. Blanchard, 1895; (B) Histological view of metacercariae of *Opisthorchis* sp. adhered on the gonads of *M. amazonicum*; (C) Metacercariae of *Opisthorchis* sp. in highlighted view; (D) Schematic drawing of the metacercaria of *Opisthorchis* sp. indicating the oral sucker (OS), ventral sucker (VS) and brownish pigment granules and an O-shaped excretory bladder (EB). Scale bars (A) = 0.5 cm; (C, D) = 100 µm.

From North America, many studies reported the presence of trematodes in impacted environments; it indicates the transmission through different routes such as intestinal transport due to tourist flow, fresh oysters exports for human consumption, aquarium trade with parasitised fishes, introduction of exotic molluscs in subtropical and tropical areas (e.g. *Lissachatina fulica* (Bowdich, 1822) and *M. tuberculata*) and climate changes setting new migratory routes of birds and fishes transporting the parasites into new regions (MURRAY; HAINES, 1969; NOLLEN; MURRAY, 1978; ISMAIL; ARIF, 1991; SCHOLZ; SALGADO-MALDONADO, 2000; DÍAZ; HERNANDEZ; BASHIRULLAH, 2002; TOLLEY-JORDAN; OWEN, 2008; PINTO; MELO, 2012).

In Brazil, the three main trematode parasite into Opisthorchiidae observed for the central and southeastern regions are *Centrocestus formosanus*, *Opisthorchis felineus* (Rivolta, 1884) and *Opisthorchis viverrini* (Poirier, 1886), with reports as endoparasites in snails (intermediate host), cats and potentially in humans (definitive hosts), as observed in case studies reported from Asian regions (OLIVEIRA et al., 2005).

Additionally, in Brazil, the trematode species of the family Heterophyidae are widely reported as intermediate hosts of the exotic species *M. tuberculata* (in metacercariae stages), with reports of parasitism in the states of Rio de Janeiro by Thiengo et al. (2001), Boaventura et al. (2002) and Bogéa; Cordeiro; Gouveia (2005); Bahia by Lemos et al. (2007); Minas Gerais by Carneiro et al. (2004) and Pinto et al. (2005); and in Brasília by Andrade et al. (2008). Additionally, the pleurolofocerca stage was reported as an endoparasite in birds, fishes and in the biliary system of humans (PINTO; MATI; MELO, 2013).

Some authors indicate the mollusc *M. tuberculata* as specific host of the families Philophthalmidae Looss, 1899 and Echinochasmidae Odhner, 1910 from Asian region (SUKONTASON et al., 2005; SAYASONE et al., 2009). However, these studies report only molluscs as the main intermediate hosts, with no observations in other aquatic invertebrates such as crustaceans, as observed herein.

In Brazilian clinical cases of trematodes as human parasites (definitive host), there are reports of *Fasciola hepatica* Linnaeus, 1758 from states of Rio de Janeiro, São Paulo and Minas Gerais, with parasites occupying the hepatic and biliary areas, acquired by eating fish from contaminated regions; additionally, Chieffi et al. (1990) observed a human case of parasitism of *Ascocotyle* sp. Looss, 1899 (in a 31-year-old woman) due to the ingestion of uncooked fish. Dias et al. (1992) observed the presence of *Clonorchis sinensis* (Cobbold, 1875) Looss, 1907 in Asian immigrants in Brazilian territory.

An infestation by *Paragonimus westermani* (Kerbert, 1878) Braun, 1899 was observed in the state of Bahia by Lemos et al. (2007), in one adult man, showing pulmonary infection, probably caused by consumption of uncooked molluscs and shrimps. Another potential trematodes transmitter are the molluscs of the genus *Biomphalaria* Preston, 1910 (intermediate host), which are widely reported in degraded aquatic environments, and used as intermediate hosts of a wide range of parasites (GIOVANELLI; VIEIRA; SILVA, 2003).

On other hand, another route of trematode contamination in humans, is the direct contact with the faeces of domestic animals (dogs and cats), once these domestic animals in the process of hunting, may consume fishes, birds, mice and other parasitised mammals in urbanised regions. Humans, in contact with the faeces of domestic animals, become definitive hosts.

The proliferation of trematode parasites in humans causes abdominal pain, hepatic and biliary degradation (in severe cases, bladder and bile cancer - Biliary Tract Carcinoma), diarrhoea, vomiting and respiratory problems; due to the absence of basic sanitation in many regions, especially in northern Brazil, these parasites are introduced into the aquatic environment, where they complete their cycle, spreading to other regions, seeking other intermediate / definitive hosts (COSTA; LIMA; COSTA, 1984).

In an experiment performed by Fan (1998), it was observed the high resistance of the trematode *C. sinensis* in metacercariae stages, recovered as parasite of freshwater fish *Pseudorasbora pamu* (Temminck & Schlegel, 1846): the fish was frozen for 7 days at a temperature of - 20 °C, however, the metacercariae remained alive and with high parasitic potential. Additionally, the same author analysed the resistance of *C. sinensis* in fish salting process, finding that the parasite remains active for up to 7 days in high salt concentration (above 10 grams of salt by cm² of fish body).

In observations made by Phyo Myint et al. (2020) in public supermarket from Tachileik (Myanmar), eight popular fish species parasitised by four metacercariae

species were found: *O. viverrini*, *Haplorchis taichui* (Nishigori, 1924); *Haplorchis pumilio* (Looss, 1896) and the genus *Haplorchoides* Chen, 1949; it indicates the high resistance in uncooked dishes consumed by the population (above 48 hours), emphasising the high risk of contamination of the population through the consumption of contaminated fish.

According to Fan (1998) and Aukkanimart et al. (2017) to prevent the presence of the parasites in humans, it is recommended the consumption of well-cooked fishery products, additionally being effective against the parasite, the consumption in salted and marinated forms or kept frozen below - 20°C for long periods.

In the state of Pará, the Amazon River prawn *M. amazonicum* is commonly sold fresh and dry salted in open-air markets and supermarkets and largely used in the local cuisine. The artisanal production process of dry salted prawn involves cooking in brine followed by natural drying, which can be an alternative for inactivating the parasite and should be studied more carefully in future works. The consumption of fresh shrimp may favour the occurrence of trematodes, in particular, *Opisthorchis* sp. in humans.

The high urbanisation process as observed at Belém metropolitan area (see Figure 1), associated to regions without basic sanitation and the eutrophication process of the Guamá River (one of the fishing locations of the Amazon River prawn consumed in the area), makes us alert the presence of the trematode *Opisthorchis* sp. parasitising a new intermediate host species and the potential risk of parasitism in humans.

4 Conclusions

In conclusion, we would like to alert the authorities and the population about the risk of contamination by metacercariae in humans (definitive host), due to inadequate consumption (for example, not well cooked in dishes, such as sushi, steamed or in salads) of aquatic organisms such as molluscs, crustaceans and fishes, collected in the Guamá River and adjacent waters, which may become a serious health problem in the region.

CREDIT AUTHORSHIP CONTRIBUTION STATEMENT

DEGM and FAAJ conceived the research ideas, designed the study and writing the manuscript; MGM and IHAC performed the first draft of this manuscript and revisions along the main text.

DECLARATION OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence this study.

FUNDING SOURCE

No financial contribution was used for the development of this article.

ACKNOWLEDGEMENTS

The authors would like to thank the *Centro Nacional de Pesquisa e Conservação da Biodiversidade Marinha do Norte - CEPNOR / ICMBio* for the support and laboratory for this study. Additionally, the authors would like to thank the anonymous reviewers for their valuable comments throughout this paper.

REFERENCES

ANDRADE, C.P.; PINTO, H.A.; COSCARELLI, D.; VIDIGAL, T.H.D.A.; De MELO, A.L. Infecção natural de *Melanoides tuberculatus* (Caenogastropoda: Thiaridae) por larvas de trematódeos digenéticos em Brasília, Distrito Federal. In: CONGRESSO BRASILEIRO DE ZOOLOGIA, 27., 2008, Curitiba. Resumos [...]. Curitiba: Sociedade Brasileira de Zoologia, 2008. p. 1483. Available from: <https://acervodigital.ufpr.br/handle/1884/34103>. Accessed on: 12 Jul. 2023.

AUKKANIMART, R.; BOONMARS, T.; SRIRAJ, P.; SRIPAN, P.; SONGSRI, J.; RATANASUWAN, P.; LAUMMAUNWAI, P.; SUWANANTRAI, A.; AUNPROMMA, S.; KHUEANGCHAINGKHWANG, S.; PUMHIRUNROJ, B.; ARTCHAYASAWAT, A.; KHUNTIKEO, N.; LOILOME, W.; NAMWAT, N.; YONGVANIT, P.; BOONJARASPINO, S. Carcinogenic liver fluke and others contaminated in pickled fish of Northeastern Thailand. *Asian Pacific Journal of Cancer Prevention*, v. 18, n. 2, p. 529-533, 2017. Available from: <https://doi.org/10.22034/APJCP.2017.18.2.529>.

BELIZARIO, J.V.; BERSABE, M.J.; HILOMEN, V.Y.; PALLER, G.V.; BUGAYON, M.G. Intestinal heterophyiasis: an emerging food-borne parasitic zoonosis in southern Philippines. *The Southeast Asian Journal of Tropical Medicine and Public Health*, v. 32, p. 36-42, 2001. Available from: <https://pubmed.ncbi.nlm.nih.gov/12041601>. Accessed on: 6 Jul. 2023.

BLANCHARD, R.A.E. Animaux parasites [Preliminary notice]. *Bulletin de la Société Zoologique de France*, v. 20, n. 8-9, p. 217, 1895. Available from: <https://www.marinespecies.org/aphia.php?p=sourcedata&ils&id=404892>. Accessed on: 04 Jul. 2023.

BOAVENTURA, M.F.; FERNANDEZ, M.A.; THIENGO, S.C.; SILVA, R.E.; MELO, A.L. Formas larvais de Trematoda provenientes de gastrópodes límnicos da microrregião Rio de Janeiro, sudeste do Brasil. *Lundiana*, V. 3, 45-49, 2002. Available from: <https://doi.org/10.35699/2675-5327.2002.21791>.

BOGÉA, T.; CORDEIRO, F.M.; GOUVEIA, J.S. *Melanoides tuberculatus* (Gastropoda: Thiariidae) as intermediate host of Heterophyidae (Trematoda: Digenea) in Rio de Janeiro metropolitan area, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo*, v. 47, p. 87-90, 2005. Available from: <https://doi.org/10.1590/S0036-46652005000200005>.

BOWDICH, T.E. *Elements of conchology, including the fossil genera and the animals. Part I. Univalves*. Paris: J. Smith, 1822. Available from: <https://doi.org/10.5962/bhl.title.77504>.

CARNEIRO, A.C.A.V.; SANTOS, V.C.; SOUSA, F.B.C.; MELO, A.L. Infecção natural de *Melanoides tuberculatus* (Müller, 1774), na represa da Pampulha, Belo Horizonte, MG. In: CONGRESSO BRASILEIRO DE ZOOLOGIA, 25., 2004, Brasília. Resumos [...] Brasília: Sociedade Brasileira de Zoologia, 2004. p. 292.

CHAI, J.Y.; SOHN, W.M.; NA, B.K.; PARK, J.B.; JEOUNG, H.G.; HOANG, E.H.; HTOON, T.T.; TIN, H.H. Zoonotic trematode metacercariae in fish from Yangon, Myanmar and their adults recovered from experimental animals. *Korean Journal of Parasitology*, v. 55, n. 6, p. 631-641, 2017. Available from: <https://doi.org/10.3347/kjp.2017.55.6.631>.

CHEN, H.T. Systematic consideration of some heterophyid trematodes in the subfamilies Haplorchinae and Stellantchasminae. *Annals of Tropical Medicine & Parasitology*, v. 43 (3-4), 304-312, 1949. Available from: <https://doi.org/10.1080/00034983.1949.11685415>.

CHIEFFI, P.; LEITE, O.H.; DIAS, R.M.D.S.; TORRES, D.M.A.V.; MANGINI, A.C.S. Human parasitism by *Phagicola* sp. (Trematoda, Heterophyidae) in São Paulo State, Brazil. *Revista do Instituto de Medicina Tropical*, v. 32, n. 4, p. 285-288, 1990. Available from: <https://doi.org/10.1590/S003646651990000400008>.

COBBOLD, T.S. The new human fluke. *Lancet*, v. 2716, n. 2, p. 1-423, 1875. Available from: [https://doi.org/10.1016/S0140-6736\(02\)48496-6](https://doi.org/10.1016/S0140-6736(02)48496-6).

PRESENCE OF METACERCARIAE OF *Opisthorchis* sp. BLANCHARD, 1895 (TREMATODA: OPISTHORCHIIDAE) IN NEW HOST *Macrobrachium amazonicum* (HELLER, 1862), FROM THE NORTHERN BRAZIL: AN EMERGING RISK TO HUMANS BY THE SHRIMP CONSUMPTION

COSTA, H.M.A.; LIMA, W.S.; COSTA, J.O. *Phagicola arnaldoi* (Travassos, 1928) Travassos, 1929 (Trematoda, Heterophyidae) em *Canis familiaris*. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, v. 36, p. 591-595, 1984. Available from: <https://www.scielo.br/j/abmvz/>. Accessed on: 03 Aug. 2023.

COSTA, G.M.M.; RIBEIRO, H.M.C.; PANTOJA, D.N.S.M. Classificação do nível de trofia de um trecho do Rio Guamá, Belém - Pará. *Revista Geoaraguaia*, v. 12, n. 1, p. 176-191, 2022. Available from: <https://periodicoscientificos.ufmt.br/ojs/index.php/geo/article/view/11681>. Accessed on: 11 July 2023.

DIAS, R.M.D.S.; MANGINI, A.C.S.; TORRES, D.M.A.G.V.; VELLOSA, S.A.G.; SILVA, R.M.; SILVA, M.I.P.G. Introdução de *Clonorchis sinensis* por imigrantes do leste asiático no Brasil e a suspensão da obrigatoriedade de exams laboratoriais para obtenção de vistos de permanência. *Revista Brasileira de Análises Clínicas*, v. 24, p. 29-30, 1992. Available from: <https://pesquisa.bvsalud.org/portal/resource/pt/lil-119549>. Accessed on: 18 Jul. 2023.

DÍAZ, M.T.; HERNANDEZ, L.E.; BASHIRULLAH, A.K. Experimental life cycle of *Philophthalmus gralli* (Trematoda: Philophthalmidae) in Venezuela. *Revista de Biología Tropical*, v. 50, p. 629-641, 2002. Available from: <https://revistas.ucr.ac.cr/index.php/rbt/article/view/16587>. Accessed on: 16 Jul. 2023.

DAO, H.T.; DERMAUW, V.; GABRIEL, S.; SUWANNATRAI, A.; TESANA, S.; NGUYEN, G.T.; DORNY, P. *Opisthorchis viverrini* infection in the snail and fish intermediate hosts in Central Vietnam. *Acta Tropical*, v. 170, p. 120-125, 2017. Available from: <https://doi.org/10.1016/j.actatropica.2017.02.028>.

EMPARANZA, E.; ULLOA, R.; MONTIEL-RAMOS, A.; MOLINA-OCAMPO, R. First record of the association of the crab *Pinnaxodes gigas* (Decapoda: Pinnotheridae) with the geoduck clam *Panopea globosa* (Bivalvia: Hiatellidae). *Marine Biodiversity Records*, v. 4, p. e40(1-3), 2011. Available from: <https://doi.org/10.1017/S1755267211000352>.

FAN, P.C. Viability of metacercariae of *Clonorchis sinensis* in frozen or salted freshwater fish. *International Journal for Parasitology*, v. 28, n. 4, p. 603-605, 1998. Available from: [https://doi.org/10.1016/s0020-7519\(97\)00215-4](https://doi.org/10.1016/s0020-7519(97)00215-4).

GIOVANELLI, A.; VIEIRA, M.V.; SILVA, C.L.P.A.C. Apparent competition through facilitation between *Melanoides tuberculata* and *Biomphalaria glabrata* and the control of schistosomiasis. *Memórias do Instituto Oswaldo Cruz*, v. 98, n. 3, p. 429-431, 2003. Available from: <https://doi.org/10.1590/S0074-02762003000300025>.

GÓNGORA-GÓMEZ, A.M.; MUÑOZ-SEVILLA, N.P.; HERNÁNDEZ SEPÚLVEDA, J.A.; GARCIA-ULLOA, M. Association between the pen shell *Atrina tuberculosa* and the shrimp *Pontonia margarita*. *Symbiosis*, v. 66, p. 107-110, 2015. Available from: <https://doi.org/10.1007/s13199-015-0342-2>.

HELLER, C. Beiträge zur näheren Kenntnis der Macrouren. *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften*, v. 1, n. 5, p. 389-426, 1862. Available from: <https://www.biodiversitylibrary.org/item/34882#page/7/mode/1up>. Accessed on: 15 Jun. 2023.

HOP, N.T.; DE, N.V.; MURRELL, D.; DALSGAARD, A. Occurrence and species distribution of fishborne zoonotic trematodes in wastewater-fed aquaculture in northern Vietnam. *Tropical Medicine & International Health*, v. 12, n. s2, p. 66-72, 2007. Available from: <https://doi.org/10.1111/j.1365-3156.2007.01943.x>.

ISMAIL, N.S.; ARIF, A.M.S. Larval trematodes of *Melanoides tuberculatus* (Müller, 1774) (Gastropoda: Prosobranchia) in brackish spring, United Arab Emirates. *Japanese Journal of Parasitology*, v. 40, p. 157-169, 1991. Available from: <https://doi.org/10.1007/BF00013997>.

JEON, H.K.; LEE, D.; PARK, H.; MIN, D.Y.; RIM, H.J.; ZHANG, H.; YANG, Y.; LI, X.; EON, K.S. Human infections with liver and minute intestinal flukes in Guangxi, China: Analysis by DNA sequencing, ultrasonography, and immunoaffinity chromatography. *Korean Journal of Parasitology*, v. 50, n. 4, p. 391-394, 2012. Available from: <https://doi.org/10.3347/kjp.2012.50.4.391>.

KERBERT, C. Zur Trematoden-kenntnis. *Zoologischer Anzeiger*, v. 1, p. 271-273, 1878. Available from: <https://www.biodiversitylibrary.org/page/30123473#page/285/mode/1up>. Accessed on: 22 July 2023.

LEMOS, A.C.M.; COELHO, J.C.; MATOS, E.D.; MONTAL, G.; AGUIAR, F.; BADARÓ, R. Paragonimiasis: first case reported in Brazil. *Brazilian Journal of Infectious Diseases*, v. 11, p. 153-156, 2007. Available from: <https://doi.org/10.1590/S1413-86702007000100031>.

PRESENCE OF METACERCARIAE OF *Opisthorchis* sp. BLANCHARD, 1895 (TREMATODA: OPISTHORCHIIDAE) IN NEW HOST *Macrobrachium amazonicum* (HELLER, 1862), FROM THE NORTHERN BRAZIL: AN EMERGING RISK TO HUMANS BY THE SHRIMP CONSUMPTION

LINNAEUS, C. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis.* Editio decima, reformata [10th revised edition], Laurentius Salvius: Holmiae, V. 1, 824 p., 1758. Available from: <https://www.biodiversitylibrary.org/item/10325#page/2/mode/1up>. Accessed on: 18 Jul. 2023.

LOOSS, A. Weitere Beiträge zur Kenntniss der Trematoden-Fauna Aegyptens, zugleich Versuch einer natürlichen Gliederung des Genus *Distomum* Retzius. *Zoologische Jahrbücher*, V. 12, 521-784, 1899. Available from: <https://www.biodiversitylibrary.org/page/10220635#page/531/mode/1up>. Accessed on: 22 July 2023.

LOOSS, A. Recherches sur la faune parasitaire de l'Égypte. Première partie. *Mémoires de l'Institut Egyptien. Le Caire*, V. 3, 1-252, 1896. Available from: <https://doi.org/10.11588/diglit.12251#0013>.

LOOSS, A. On some parasites in the Museum of the School of Tropical Medicine, Liverpool. With a contribution on a case of distomiasis of the liver and the rectum, by Edward Cuffey. *Annals of Tropical Medicine and Parasitology*, v. 1, n. 1, p. 121-154, 1907. Available from: <https://doi.org/10.1080/00034983.1907.11719254>.

LOVIS, L.; MAK, T.K.; PHONGLUXA, K.; SOUKHATHAMMAMONG, P.; SAYASONE, S.; AKKHAVONG, K.; ODERMATT, P.; KEISER, J.; FELGER, I. PCR diagnosis of *Opisthorchis viverrini* and *Haplorchis taichui* infections in a Lao community in an area of endemicity and comparison of diagnostic methods for parasitological field surveys. *Journal of Clinical Microbiology*, v. 47, p. 1517-1523, 2009. Available from: <https://doi.org/10.1128/JCM.02011-08>.

MATTOS, D.P.B.G.; VERICIMO, M.A.; SÃO CLEMENTE, S.C. O pescado e os cestóides Trypanorhyncha - do aspecto higiênico ao potencial alergênico. *Veterinária Notícias*, v. 19, n. 2, p. 127-139, 2013. Available from: <https://seer.ufu.br/index.php/vetnot/article/view/2334>. Accessed on: 22 July 2023.

MÜLLER, O.F. *Vermium terrestrium et fluviatilium, seu animalium infusorium, Helminthicorum, et testaceorum, non marinorum, succincta historia.* vol 2: I-XXXVI. Havniae et Lipsiae, apud Heineck et Faber, ex officina Molleriana, 214 p., (10 unnumbered pages), 1774. Available from: <https://doi.org/10.5962/bhl.title.46299>.

MURRAY, H.D.; HAINES, D. *Philophthalmus* sp. (Trematoda) in *Tarebia granifera* and *Melanoides tuberculatus* in south Texas. *Bulletin of the American Malacological Union*, p. 44-45, 1969. Available from: <https://www.biodiversitylibrary.org/bibliography/124945>. Accessed on: 08 Jul. 2023.

NISHIGORI, M. The life cycles of two new species of Heterophyidae, *Monorchotrema taihoku* and *M. taichui*, found in Formosa. Preliminary note. *Taiwan Igakkai Zasshi, Taihoku*, V. 237, 567-570, 1924. Available from: <https://doi.org/10.2307/3271705>.

NOLLEN, P.M.; MURRAY, H.D. *Philophthalmus gralli*: identification, growth characteristics, and treatment of an oriental eyeworm of birds introduced into the continental United States. *Journal of Parasitology*, v. 94, n. 1, p. 178-180, 1978. Available from: <https://doi.org/10.2307/3279646>.

NOUDEH, M.A.; PAZOOKI, J. The first record of *Ascocotyle (Phagicola) longa* (Digenea: Heterophyidae) metacercariae as the fish-borne zoonotic trematode in *Chelon auratus* and *Chelon saliens* (Pisces: Mugilidae) around the cage culture in the Caspian Sea, Iranian. *Journal of Parasitology*, v. 65, n. 4, p. 365-370, 2019. Available from: <https://doi.org/10.17420/ap6504.222>.

ODHNER, T. Nordostafrikanische Trematoden, grösstenteils vom Weissen Nil (von der schweidischen zoologischen Expedition gesammelt). In: JÄGERSKIÖLD, L.A. (Ed.) *Results of the Swedish zoological expedition to Egypt and the White Nile, 1901, under the direction of L.A. Jägerskiöld*. Uppsala: Library of the Royal University of Uppsala, 1910. v. 23A, p. 1-170. Available from: <https://www.biodiversitylibrary.org/item/62150#page/833/mode/1up>. Accessed on: 14 Jul. 2023.

OLIVEIRA, P.M.A.; PIRES, P.R.R.; GINJA, M.J.; PIRES, I.; CARDOSO, L.; ANTUNES, L.; RODRIGUES, M. *Opisthorchis felineus* in cat: case report. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, v. 57, n. 4, p. 556-558, 2005. Available from: <https://doi.org/10.1590/S0102-09352005000400020>.

PFÄFFLE, M.; LITWIN, N.; MUDERS, S.V.; PETNEY, T. N. The ecology of tick-borne diseases. *International Journal of Parasitology*, v. 43, n. 12-13, p. 1059-1077, 2013. Available from: <https://doi.org/10.1016/j.ijpara.2013.06.009>.

PHYO MYINT, E.E.; SEREEMASPUN, A.; ROCKLÖV, J.; NITHIKATHKUL, C. Discovery of Carcinogenic Liver Fluke Metacercariae in Second Intermediate Hosts and Surveillance on Fish-Borne Trematode Metacercariae Infections in Mekong Region of Myanmar. *International Journal of Environmental Research and Public Health*, v. 17, n. 11, article 4108, 2020. Available from: <https://doi.org/10.3390/ijerph17114108>.

PRESENCE OF METACERCARIAE OF *Opisthorchis* sp. BLANCHARD, 1895 (TREMATODA: OPISTHORCHIIDAE) IN NEW HOST *Macrobrachium amazonicum* (HELLER, 1862), FROM THE NORTHERN BRAZIL: AN EMERGING RISK TO HUMANS BY THE SHRIMP CONSUMPTION

PINTO, R.M.; SANTOS, L.C.; TORTELLY, R.; MENEZES, R. C.; MORAES, W.; JUVENAL, J.C.; GOMES, D.C. Pathology and first report of natural infections of the eye trematode *Philophthalmus lachrymosus* Braun, 1902 (Digenea, Philophthalmidae) in a non-human mammalian host. *Memórias do Instituto Oswaldo Cruz*, v. 100, 579-583, 2005. Available from: <https://doi.org/10.1590/S0074-02762005000600012>.

PINTO, H. A.; MELO, A. L. *Melanoides tuberculata* (Mollusca: Thiaridae) as an intermediate host of *Centrocestus formosanus* (Trematoda: Heterophyidae) in Brazil. *Revista do Instituto de Medicina Tropical de São Paulo*, v. 52 (4), 207-210, 2010. Available from: <https://doi.org/10.1590/S0036-46652010000400008>.

PINTO, H. A.; MELO, A. L. Metacercariae of *Centrocestus formosanus* (Trematoda: Heterophyidae) in *Australoheros facetus* (Pisces: Cichlidae) in Brazil. *Revista Brasileira de Parasitologia Veterinária*, v. 21 (3), 334-337, 2012. Available from: <https://doi.org/10.1590/S1984-29612012000300032>.

PINTO, H. A.; MATI, V. L. T.; MELO, A. L. New records and a checklist of trematodes from *Butorides striata* (Aves: Ardeidae). *Revista Mexicana de Biodiversidad*, v. 84 (4), 1100-1110, 2013. Available from: <https://doi.org/10.7550/rmb.34431>.

PRESTON, H. B. Additions to the non-marine molluscan fauna of British and German East Africa and Lake Albert Edward. *The Annals and Magazine of Natural History*, v. 6, 526-536, 1910. Available from: <https://www.biodiversitylibrary.org/page/15629297#page/556/mode/1up>. Accessed on: 15 Jul. 2023.

POIRIER, J. Trématodes nouveaux ou peu connus. *Bulletin de la Société Philomathique*, Paris, v. 7 (10), 20-40, 1886.

PUMIDONMING, W.; KATAHIRA, H.; IGARASHI, M.; SALMAN, D.; ABDELBASET, A. E.; SANGKAO, K. Potential risk of a liver fluke *Opisthorchis viverrini* infection brought by immigrants from prevalent areas: A case study in the lower Northern Thailand. *Acta Tropical*, v. 178, 213-218, 2018. Available from: <https://doi.org/10.1016/j.actatropica.2017.11.023>.

RIVOLTA, S. Sopra una specie di Distomanelgatto e nel cane. *Giornale di anatomia, fisiologia e patologia degli animali*, v. 10 (1), 20-28, 1884.

SAYASONE, S.; TESANA, S.; UTZINGER, J.; HATZ, C.; AKKHAVONG, K.; ODERMATT, P. Rare human infection with the trematode *Echinocasmus japonicus* in Lao PDR. *Parasitology International*, v. 58, 106-109, 2009. Available from: <https://doi.org/10.1016/j.parint.2008.11.001>.

SITHITHAWORN, P.; ANDREWS, R. H.; VAN DE, N.; WONGSAROJ, T.; SINUON, M.; ODERMATT, P.; NAWA, Y.; LIANG, S.; BRINDLEY, P. J.; SRIPA, B. The current status of opisthorchiasis and clonorchiasis in the Mekong Basin. *Parasitology International*, v. 61, 10-16, 2012. Available from: <https://doi.org/10.1016/j.parint.2011.08.014>.

SCHOLZ, T.; DITRICH, O.; GIBODA, M. Study on the surface morphology of the developmental stages of the liver fluke, *Opisthorchis viverrini* (Trematoda: Opisthorchiidae). *Annales de parasitologie humaine et comparée*, v. 67 (3), 82-90, 1992. Available from: <https://doi.org/10.1051/parasite/199267382>.

SCHOLZ, T.; SALGADO-MALDONADO, G. The introduction and dispersal of *Centrocestus formosanus* (Nishigori, 1924) (Digenea: Heterophyidae) in Mexico: a review. *American Midland Naturalist*, v. 143, 185-200, 2000. Available from: [https://doi.org/10.1674/00030031\(2000\)143\[0185:TIADOC\]2.0.CO;2](https://doi.org/10.1674/00030031(2000)143[0185:TIADOC]2.0.CO;2).

SUKONTASON, K.; UNPUNYO, P.; SUKONTASON, K. L.; PIANGJAI, S. Evidence of *Haplorchis taichui* infection as pathogenic parasite: three case reports. *Scandinavian Journal of Infectious Diseases*, v. 37, 388-390, 2005. Available from: <https://doi.org/10.1080/00365540510034473>.

THATCHER, V. E. Trematódeos neotropicais. Manaus: Instituto Nacional de Pesquisas da Amazônia, 553p., 1993. Available from: https://openlibrary.org/books/OL318092M/Tremato%CC%81deos_neotropicais. Accessed on: 02 Jul. 2023.

THIENGO, S. C.; FERNANDEZ, M. A.; BOAVENTURA, M. F.; GRAULT, C. E.; SILVA H. F. R.; MATTOS, A. C.; SANTOS, S. B. Freshwater snails and Schistosomiasis mansoni in the state of Rio de Janeiro, Brazil: I - Metropolitan Mesoregion. *Memórias do Instituto Oswaldo Cruz*, v. 96 (suppl. 1), 177-184, 2001. Available from: <https://doi.org/10.1590/S0074-02762002000900010>.

TEMMINCK, C. J.; SCHLEGEL, H. Pisces. In: SIEBOLD, P. F. (Ed.): *Fauna Japonica, sive descriptio animalium, quae in itinere per Japoniam. suscepto annis 1823-1830 collegit, notis, observationibus et adumbrationibus illustravit Ph. Fr. de Siebold*. Lugduni Batavorum [Leiden] (A. Arnz et soc.). Parts 10-14: 173-269, 1846. Available from: <https://www.biodiversitylibrary.org/item/217507#page/41/mode/1up>. Accessed on: 15 Jul. 2023.

TOLLEY-JORDAN, L. R.; OWEN, J. M. Habitat influences snail community structure and trematode infection levels in a spring-fed river, Texas, USA. *Hydrobiologia*, v. 600, 29-40, 2008. Available from: <https://doi.org/10.1007/s10750-007-9173-3>.

PRESENCE OF METACERCARIAE OF *Opisthorchis* sp. BLANCHARD, 1895 (TREMATODA: OPISTHORCHIIDAE) IN NEW HOST *Macrobrachium amazonicum* (HELLER, 1862), FROM THE NORTHERN BRAZIL: AN EMERGING RISK TO HUMANS BY THE SHRIMP CONSUMPTION

TOUCH, S.; KOMALAMISRA, C.; RADOMYOS, P.; WAIKAGUL, J. Discovery of *Opisthorchis viverrin* metacercariae in freshwater fish in southern Cambodia. *Acta Tropica*, V. 111, 108-113, 2009. Available from: <https://doi.org/10.1016/j.actatropica.2009.03.002>.

VAJRASTHIRA, S.; HARINASUTA, C.; KOMIYA, Y. The Morphology of the Metacercaria of *Opisthorchis Viverrini*, with Special Reference to the Excretory System. *Annals of Tropical Medicine & Parasitology*, V. 55 (4), 413-418, 1961. Available from: <https://doi.org/10.1080/00034983.1961.11686068>.

WoRMS (2023). *Opisthorchis* Blanchard, 1895. Available from: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=108622>. Accessed on: 02 Jul. 2023.